



GB04/04770



INVESTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road

Newport

South Wales  
NP10 8QQ

RECD 07 DEC 2004

WIPO

PC

**PRIORITY DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 22 November 2004

BEST AVAILABLE COPY

Patents Form 1/77

Patents Act 1977  
(Rule 16)THE  
Patent  
Office

1/77

## Request for grant of a patent

*(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form.)*THE PATENT OFFICE  
PE

04 DEC 2003

RECEIVED BY FAX

Fee: £0

The Patent Office

Cardiff Road  
Newport  
Gwent NP9 1RH

1. Your reference

46978-GB01/JMP EC03 E857006-1 001631

77700 0-00-0328060.9

2. Patent application number  
*(The Patent Office will fill in this part)*

0328060.9

3. Full name, address and postcode of the or of  
each applicant *(underline all surnames)*SOCIETE DE CONSEILS DE RECHERCHES ET  
D'APPLICATIONS SCIENTIFIQUES (S.C.R.A.S)  
42, rue du Docteur Blanche  
75016 Paris  
FRANCEPatents ADP number *(if you know it)*

6199020002

If the applicant is a corporate body, give the  
country/state of incorporation

France

4. Title of the invention

Botulinum Toxin Treatment

5. Full name, address and postcode in the United  
Kingdom to which all correspondence relating  
to this form and translation should be sentReddie & Grose  
16 Theobalds Road  
LONDON  
WC1X 8PLPatents ADP number *(if you know it)*

91001

6. If you are declaring priority from one or more  
earlier patent applications, give the country  
and the date of filing of the or of each of these  
earlier applications and *(if you know it)* the or  
each application number

Country

Priority application  
*(if you know it)*Date of filing  
*(day/month/year)*7. If this application is divided or otherwise  
derived from an earlier UK application,  
give the number and the filing date of  
the earlier application

Number of earlier application

Date of filing  
*(day/month/year)*8. Is a statement of inventorship and of right  
to grant of a patent required in support of  
this request? *(Answer 'Yes' if:*

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an  
applicant, or
- c) any named applicant is a corporate body.  
*See note (d))*

YES

Patents Form 1/77

0087929 04 Dec 03 01:10

**Patents Form 1/77**

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document.

Continuation sheets of this form

Description	12
Claim(s)	2
Abstract	0
Drawing(s)	0

10. If you are also filing any of the following, state how many against each item.

Priority documents	0
Translations of priority documents	0
Statement of inventorship and right to grant of a patent ( <i>Patents Form 7/77</i> )	0
Request for preliminary examination and search ( <i>Patents Form 9/77</i> )	1
Request for substantive examination ( <i>Patents Form 10/77</i> )	0
Any other documents (please specify)	

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date

4 December 2003

*Reddie & Grose*

12. Name and daytime telephone number of person to contact in the United Kingdom

J M DAVIES  
01223 360350

**Warning**

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or such direction has been revoked.

**Notes**

1. If you are filing a patent application in the United Kingdom, you must also file a copy of the application with the Patent Office in the United Kingdom.

2. If you are filing a patent application in the United Kingdom, you must also file a copy of the application with the Patent Office in the United Kingdom.

3. If you are filing a patent application in the United Kingdom, you must also file a copy of the application with the Patent Office in the United Kingdom.

4. If you are filing a patent application in the United Kingdom, you must also file a copy of the application with the Patent Office in the United Kingdom.

5. If you are filing a patent application in the United Kingdom, you must also file a copy of the application with the Patent Office in the United Kingdom.

6. If you are filing a patent application in the United Kingdom, you must also file a copy of the application with the Patent Office in the United Kingdom.

7. If you are filing a patent application in the United Kingdom, you must also file a copy of the application with the Patent Office in the United Kingdom.

**DUPLICATE****Botulinum Toxin Treatment**

The present invention relates to the use of botulinum toxin for the preparation of a medicament intended to topically treat urinary disorders and notably spastic bladder.

The anaerobic, gram positive bacterium *Clostridium botulinum* produces a potent polypeptide neurotoxin, botulinum toxin, which causes a neuromuscular illness in humans and animals referred to as botulism. The spores of *Clostridium botulinum* are found in soil and can grow in improperly sterilized and sealed food tins, which are the cause of many of the cases of botulism. The effects of botulism typically appear 18 to 36 hours after eating the foodstuffs infected with a *Clostridium botulinum* culture or spores. The botulinum toxin can apparently pass unattenuated through the lining of the gut and attack peripheral motor neurons. Symptoms of botulinum toxin intoxication can progress from difficulty walking, swallowing, and speaking to paralysis of the respiratory muscles and death.

Botulinum toxin type A is the most lethal natural biological agent known to man. About 50 picograms of a commercially available botulinum toxin type A (purified neurotoxin complex) correspond to the LD<sub>50</sub> in mice.

Seven immunologically distinct botulinum neurotoxins have been characterized, these being respectively botulinum neurotoxin serotypes A, B, C<sub>1</sub>, D, E, F and G each of which is distinguished by neutralization with type-specific antibodies. The different serotypes of botulinum toxin vary in the animal species that they affect and in the severity and duration of the paralysis they evoke. For example, it has been determined that botulinum toxin type A is 500 times more potent, as measured by the LD<sub>50</sub> in mice, than botulinum toxin type B. Additionally, botulinum toxin type B has been determined to be non-toxic in primates at a dose of 480 U/kg which is about 12 times the primate LD<sub>50</sub> for botulinum toxin type A. Botulinum toxin apparently binds with high affinity to cholinergic motor neurons, is translocated into the neuron and blocks the release of acetylcholine.

Although all the botulinum toxin serotypes apparently inhibit release of the neurotransmitter acetylcholine at the neuromuscular junction, they do so by affecting different neurosecretory proteins and/or cleaving these proteins at different sites. For example, botulinum types A and E both cleave the 25 kilodalton (kD) synaptosomal associated protein (SNAP-25), but they target different amino acid sequences within this protein. Botulinum toxin types B, D, F and G act on vesicle-associated protein (VAMP, also called synaptobrevin), with each serotype cleaving the protein at a

- 2 -

different site. Finally, botulinum toxin type C<sub>1</sub> has been shown to cleave both syntaxin and SNAP-25. These differences in mechanism of action may affect the relative potency and/or duration of action of the various botulinum toxin serotypes.

5 The molecular weight of the botulinum toxin protein molecule, for all seven of the known botulinum toxin serotypes, is about 150 kD. Interestingly, the botulinum toxins are released by Clostridial bacterium as complexes comprising the 150 kD botulinum toxin protein molecule along with associated non-toxin proteins. Thus, the botulinum toxin type A complex can be produced by Clostridial bacterium as 900 kD, 500 kD and 300 kD forms. Botulinum toxin types B and C<sub>1</sub> are apparently produced as only a 500  
10 kD complex. Botulinum toxin type D is produced as both 300 ka and 500 kD complexes. Finally, botulinum toxin types E and F are produced as only approximately 300 kD complexes. The complexes (i.e. molecular weight greater than about 150kD) are believed to contain a non-toxin hemagglutinin protein and a nontoxin and non-toxic nonhemagglutinin protein. These two non-toxin proteins (which along with the  
15 botulinum toxin molecule comprise the relevant neurotoxin complex) may act to provide stability against denaturation to the botulinum toxin molecule and protection against digestive acids when toxin is ingested. Additionally, it is possible that the larger (greater than about 150 kD molecular weight) botulinum toxin complexes may result in a slower rate of diffusion of the botulinum toxin away from a site of intramuscular  
20 injection of a botulinum toxin complex.

Botulinum toxin type A can be obtained by establishing and growing cultures of Clostridium botulinum in a fermenter and then harvesting and purifying the fermented mixture in accordance with known procedures. All the botulinum toxin serotypes are initially synthesized as inactive single chain proteins which must be cleaved or nicked  
25 by proteases to become neuroactive. The bacterial strains that make botulinum toxin serotypes A and G possess endogenous proteases and serotypes A and G can therefore be recovered from bacterial cultures in predominantly their active form. In contrast, botulinum toxin serotypes C<sub>1</sub>, D and E are synthesized by nonproteolytic strains and are therefore typically unactivated when recovered from culture. Serotypes B and F are  
30 produced by both proteolytic and nonproteolytic strains and therefore can be recovered in either the active or inactive form. However, even the proteolytic strains that produce, for example, the botulinum toxin type B serotype only cleave a portion of the toxin produced. The exact proportion of active to produced toxin varies depending on the strain.

- molecules in a clinical preparation will contribute to the overall protein load of the preparation, which has been linked to increased antigenicity, without contributing to its clinical efficacy. Additionally, it is known that botulinum toxin type B has, upon intramuscular injection, a shorter duration of activity and is also less potent than
- 5 botulinum toxin type A at the same dose level. High quality crystalline botulinum toxin type A can be produced from the Hall A strain of *Clostridium botulinum* with characteristics of  $3 \cdot 10^7$  U/mg, an  $A_{260}/A_{278}$  of less than 0.60 and a distinct pattern of banding on gel electrophoresis. The known Schantz process can be used to obtain crystalline botulinum toxin type A, as set forth in Schantz, E. J., et al, Properties and use
- 10 of Botulinum toxin and Other Microbial Neurotoxins in Medicine, *Microbiol Rev.* (1992), 56, 80-99. Generally, the botulinum toxin type A complex can be isolated and purified from an anaerobic fermentation by culturing *Clostridium botulinum* type A in a suitable medium. The known process can also be used, upon separation out of the non-toxin proteins, to obtain pure botulinum toxins, such as for example: purified botulinum
- 15 toxintype A with an approximately 150 kD molecular weight with a specific potency of  $1-2 \cdot 10^8$  LD<sub>50</sub> U/mg or greater, purified botulinum toxin type B with an approximately 156 kD molecular weight with a specific potency of  $1-2 \cdot 10^8$  LD<sub>50</sub> U/mg or greater, and purified botulinum toxin type F with an approximately 155 kD molecular weight with a specific potency of  $1-2 \cdot 10^7$  LD<sub>50</sub> U/mg or greater.
- 20 Already prepared and purified botulinum toxins and toxin complexes can be obtained from List Biological Laboratories, Inc. (Campbell, California, USA); the Centre for Applied Microbiology and Research (Porton Down, UK); Wako (Osaka, Japan), as well as from Sigma Chemicals (St. Louis, Missouri, USA).

- Botulinum toxin of type A was first used in man in 1981. The first therapeutic uses
- 25 related to the treatment of certain neuromuscular disorders that include blepharospasm, strabismus and hemifacial spasm. Non-type A botulinum toxin serotypes apparently have a lower potency and/or a shorter duration of activity as compared to botulinum toxin type A. Clinical effects of peripheral intramuscular botulinum toxin type A are usually seen within one week of injection. The typical duration of symptomatic relief
- 30 from a single intramuscular injection of botulinum toxin type A averages about three months.

Spastic bladder treatment by botulinum toxin was first disclosed in US patent No. 5,437,291. The method of administration disclosed is the direct injection into the bladder wall.

PCT patent application WO 99/03483 further discloses the use of botulinum toxin for the treatment of various urinary disorders. The administration of toxin consists in an injection in the lateral bladder wall by means of an endoscopic device introduced through the urethra.

5 The Applicant has now surprisingly found that it is possible to treat patients with urinary disorders and notably spastic bladder without using such an injection. This can be achieved for example by one of the following methods:

- infusing the patients' bladder with a liquid or semi-solid formulation of botulinum toxin;
- 10 - depositing a gel formulation containing botulinum toxin at the appropriate location of the patients' bladder;
- spraying a spray formulation containing botulinum toxin at the appropriate location of the patients' bladder; or
- 15 - topically applying a solid (e.g. lyophilised), semi-solid or liquid botulinum toxin formulation that is put or spread on the outer walls of a balloon which is then inflated in the bladder so as to be in contact with said bladder's wall.

By spray formulation is understood in the present application a solid or liquid formulation that is or can be split into particles or droplets with a mean diameter of less than 1 mm, preferably of less than 500  $\mu\text{m}$  and more preferably less than 100  $\mu\text{m}$ .

20 The invention therefore relates to the use of a liquid, semi-solid, gel, spray or solid (e.g. lyophilised) formulation of botulinum toxin for the preparation of a medicament intended to treat a disorder characterised by bladder spasms wherein said medicament is for administration by infusion into the bladder or local administration without injection into the bladder wall.

25 Liquid formulations may in particular be a liquid formulation that is not miscible with water and such that the formulation is less dense than water; an advantage of this execution mode is that the formulation tends to separate and preferentially be in contact with the bladder. The liquid formulations may also be water solutions (e.g. a saline water solution) or suspensions.

30 Besides, the liquid or semi-solid formulation may be replaced by a gel (including a solid gel) or another suitable formulation. This type of formulation will be  
\_\_\_\_\_

- 5 -

Alternatively, the formulation may be in the form of a spray containing the botulinum toxin. This spray may notably be created using a two-compartment capsule containing in one compartment botulinum toxin powder and in the other explosives which will bring about the spraying of botulinum toxin once explosion occurs; hence, according to  
5 this particular variant, the capsule is first brought at the appropriate location near the bladder wall and then the explosion triggered to spray the botulinum toxin on said wall.

According to a further variant of the invention, a solid (e.g. lyophilised), semi-solid or liquid botulinum toxin formulation may be coated or spread over the outer walls of a balloon. The balloon is brought into the bladder using an endoscopic or cystoscopic  
10 apparatus or, more preferably, only a catheter; it is then inflated in the bladder until it is in contact with the bladder walls. In this manner, the botulinum toxin can be delivered at the appropriate bladder wall locations, avoiding administration to the trigone. A further object of the invention is therefore the use of a solid (e.g. lyophilised), semi-solid or liquid botulinum toxin formulation spread on the outer wall of a balloon  
15 intended to be inflated inside the bladder for the preparation of a medicament intended to treat a disorder characterised by bladder spasms.

Preferably, the botulinum toxin formulation used for this invention will comprise a penetration enhancer.

To summarise, according to the invention, the formulation of botulinum toxin does not  
20 need to be injected but should simply be put in contact with the bladder wall. This mode of administration is much easier for the doctor.

According to this invention, disorders characterised by bladder spasms include notably:

- ❖ urinary incontinence due to unstable bladder or unstable detrusor sphincter;
- ❖ voiding complications due to detrusor overactivity or unstable detrusor sphincter;
- 25 ❖ detrusor sphincter dyssynergia;
- ❖ urinary retention secondary to spastic sphincter or hypertrophied bladder neck; or
- ❖ neurogenic bladder dysfunction secondary to Parkinson's disease, spinal cord injury, stroke or multiple sclerosis or characterised by a spasm reflex.

In this application, unless stated otherwise, the words "botulinum toxin" always refer to  
30 a botulinum toxin complex (made from the toxin and one or more complexing proteins) or a high purity botulinum toxin (botulinum toxin which is substantially free from any complexing protein).



By botulinum toxin complex shall be notably understood the active principles of the products Dysport<sup>®</sup> (registered trademark of the Applicant), Botox<sup>®</sup> (registered trademark of Allergan), Vistabel<sup>®</sup> (registered trademark of Allergan), Neurobloc<sup>®</sup> (registered trademark of Elan) or Myobloc<sup>®</sup> (registered trademark of Elan).

- 5 By high purity botulinum neurotoxin (type A, B, C<sub>1</sub>, D, E, F or G) is meant, in the present application, botulinum neurotoxin (type A, B, C<sub>1</sub>, D, E, F or G) free from complexes including at least another protein. In other words, a high purity botulinum neurotoxin (type A, B, C<sub>1</sub>, D, E, F or G) does not contain significant quantities of any other *Clostridium spp* derived protein than botulinum neurotoxin (type A, B, C<sub>1</sub>, D, E, F or G).
- 10

Preferably, the invention will be such that the medicament prepared is intended to treat a disorder selected from the group consisting of urinary incontinence, urinary retention and neurogenic bladder dysfunction.

- 15 A liquid or semi-solid formulation intended for bladder infusion according to the invention will preferably not have a volume exceeding 100 ml, and more preferably 80 ml. For example, a volume of 20 to 80 ml, and notably a volume of 40 to 60 ml (e.g. about 50 ml), will be appropriate. The same volumes will apply for a gel formulation intended to be deposited in the patient's bladder.

- 20 Preferably, the viscosity of the formulation according to the invention will be greater than that of the corresponding water solution thanks to the use of a thickening agent like gelatine, carboxymethyl cellulose, polyethylene glycol, glycerol, mannitol or a surfactant.

- 25 Furthermore, liquid formulations used for the invention will preferably be such that they have a viscosity from 0.5 to 500 centipoises, and more preferably from 1 to 500 centipoises. Semi-solid formulations used for the invention will preferably be such that they have a viscosity from 500 to 10,000 centipoises, and more preferably from 2,000 to 10,000 centipoises. Gel formulations used for the invention will preferably be such that they have a viscosity from 10,000 to 100,000 centipoises, and more preferably from 50,000 to 100,000 centipoises.

- 30 ~~For use in a bladder infusion the liquid or semi-solid formulation according to the~~  
~~invention will preferably have a viscosity from 0.5 to 500 centipoises, and more~~  
~~preferably from 1 to 500 centipoises. Semi-solid formulations used for the invention~~  
~~will preferably be such that they have a viscosity from 500 to 10,000 centipoises,~~  
~~and more preferably from 2,000 to 10,000 centipoises. Gel formulations used for the~~  
~~invention will preferably be such that they have a viscosity from 10,000 to 100,000~~  
~~centipoises, and more preferably from 50,000 to 100,000 centipoises.~~

- 7 -

Nevertheless, the dose of botulinum toxin to provide according to the present invention for treatment of the abovementioned diseases or disorders, varies depending on the age and body weight of the subject to be treated, as well as the state of the latter, and will be finally decided by the attending doctor or veterinarian. Such a quantity determined by the attending doctor or veterinarian is here called "therapeutically effective quantity".

For the purposes of the instant patent application, one unit (U) of botulinum toxin is defined as the LD<sub>50</sub> upon intraperitoneal injection into female Swiss Webster mice weighing 18 to 20 grams each.

Methods of treatment of disorders characterised by bladder spasms comprising the administration by infusion of a liquid, semi-solid or gel composition or spraying a spray composition according to the invention into the bladder of a patient in need thereof are also within the scope of this invention. These methods of treatment can notably be carried out under cystoscope control, but using flexible cystoscopes, mini-cystoscopes or even simple catheters will be preferred. The use of the aforesaid devices can be combined with a radioscopic control, but may also be carried out without any such control.

For the case of a spray formulation, the injection nozzle will preferably have an umbrella or conic shape in order to direct the spray in an appropriate fashion.

According to the instant invention, the infusion or contact with the botulinum toxin formulation should last any time between 10 minutes and 12 hours (overnight).

A preferred execution mode for this invention will involve the use of a balloon whose outer wall is coated with a botulinum toxin formulation, said balloon being intended to be inflated in the bladder of the patient to be treated with the purpose of releasing the toxin into the bladder wall.

The method uses a balloon that can be introduced, through the urethra, into the bladder, that can be inflated in the bladder in order to enter in contact with the bladder wall and that can also be deflated afterwards. The surface of the balloon is previously covered by a botulinum toxin formulation in order to, later, transfer said formulation as a layer between balloon and bladder wall.

The botulinum toxin formulation can be liquid, semisolid or solid and can optionally comprise one or more penetration enhancers to favour the botulinum toxin transfer from the surface to the target of the drug.

The balloon is the transfer-target device and can play an occlusive role or any other mechanical functions likely to increase botulinum toxin penetration thanks to the process and/or to the balloon surface, composition, etc.

- One of the preferred device presentations is the one used in balloon angioplasty.
- 5 However, the skilled person may contemplate any other where an inflated balloon is used in order to enter in contact with the bladder wall to ensure botulinum toxin penetration into the bladder wall.

- Hence, as for angioplasty, the device will preferably be thin and long for transluminal access to the bladder, the diameter and length being adapted to urethra and bladder. The
- 10 deflated total diameter of the balloon could be less than 1 cm and preferably less than 5 mm and the deflated total length less than 15 cm and preferably less than 10 or even 5 cm.

- The balloon device could come with a control guide wire and/or a peripheral guide catheter. For example, a thin and flexible balloon catheter can be positioned at rest in a
- 15 reservoir and when necessary expelled from the catheter (optionally through a catheter guide), leaving the balloon to expand gently into the bladder.

- One may monitor the progress of the treatment on an X-ray machine thanks to the metallic guide wire but it can also be contemplated that the wire is used as a guide to determine the appropriate position with contact on top of the bladder before treatment
- 20 application without any radiographic need, thus making the treatment even simpler to perform. The device shape and working process can be made in a way to offer contact in the target area and for instance avoiding balloon contact in the trigone or urethra area.

According to the invention method, the balloon's outer surface is to be covered with the botulinum toxin formulation to be delivered.

- 25 Different approaches with solid, semi-solid or liquid compositions of the drug can be contemplated as detailed hereafter.

- For solid formulations, one can prepare concentrated liquid aqueous or organic solutions of botulinum toxin, to wet the deflated balloon in these solutions of the drug and to dry or freeze-dry the preparation in order to obtain a solid layer of the drug on top
- 30 of the surface. Alternatively, one can dissolve or suspend the drug in a liquid and then enter in

inflammation, creating micro-lesions that will ease the transfer of the toxin through the epithelium layer. Besides, surfactant or organic solvents can be incorporated into the toxin formulation to improve the delivery. In the end, aqueous medium and/or local humidity in the bladder could rehydrate the solid composition and give the pH, concentration and liquid vehicle to transfer the toxin.

Semi-solid or gel like compositions could also be used with the balloon device. Here also one can wet the balloon surface in a botulinum toxin gel preparation or semi-solid formulation to obtain the botulinum toxin coating layer. The treatment could also be realised just by putting on top of the balloon in the guide catheter the semi-solid or gel preparation. During the process, this layer will cover the balloon's surface as a result of the balloon inflation and then be trapped, stuck and squeezed between the balloon and the bladder wall.

A liquid preparation could also be used with the device. This preparation can be already associated with the device as a layer on a deflated balloon or a liquid reservoir in the tip of the guide catheter. The liquid reservoir at the tip of the guide catheter may as well be filled using a syringe just before use with a high aqueous concentration of botulinum toxin in a small volume (e.g. 500 units of Dysport® in 0.5 ml or less of saline solution); pH modifying agents, organic solvents, viscosity and penetration enhancers may be added if necessary. This tip part will wet the balloon as it is pushed out of the guide catheter and cover it when it inflates in the bladder, allowing the toxin composition to enter in contact with the bladder wall.

Various methods of treatment can be contemplated thanks to the invention's balloon device.

For example, the balloon can be inflated and deflated several times, with a contact between the balloon's outer wall and the bladder inner wall being maintained for a certain time in between. The contact time will be chosen for optimising penetration (in general less than one hour, for instance 10 minutes) as well as the number of times the balloon will be inflated.

The pressure exerted on the bladder wall could also play a favorable plastic and penetration role. It will therefore be adapted by the treating physician to each patient. The balloon can be inflated to a relatively high pressure that can be precisely controlled (as it is normally prevented from blowing up thanks to the bladder wall). In addition, the device can be inflated or deflated several times during treatment and one can perform some pressure jerks to improve toxin penetration.

- If by chance the bladder wall gets damaged because of treatment, the healing would have the time it needs as treatments could be spaced by time intervals of 6 months or more as the toxin is effective for that duration when treating bladder disorders.

- It could also be contemplated to use an elastic catheter tipped with a balloon as a way to target local multi-injections, similarly to what is described for example in US patent No. 6,638,246. According to this variant, the balloon would be fitted with small needles (e.g. microneedles of about 1 mm length and about 130  $\mu$ m diameter) that would be attached and folded safely into the balloon's surface. As the balloon is inflated, the microneedle would penetrate the bladder wall and the botulinum toxin formulation would be injected.

- Thanks to this balloon device, the treatment dose can be accurate, already fixed and associated to the device or easily selected and prepared by the physician using standard procedures for liquid injections.

It should be mentioned that the balloon system previously described can be used for any kind of active substance that ~~can be~~ formulated as a solid, semi-solid or liquid pharmaceutical composition and that can be delivered through an epithelium surface into an accessible body reservoir or vessel like for example the bladder, the colon, the terminal ileum, the stomach, the oesophagus, the rectum, the vagina, the ear, the eye or the blood.

Possible uses of this balloon device therefore include for example:

- introducing the invention balloon coated with an anti-restenotic drug, an antithrombogenic agent, an endothelialisation promoter, a radioactive isotope, a platelet antiaggregant agent, a fibrinolytic agent, an anti-inflammatory agent, an antiproliferation agent, a radio opaque substance, a gene therapy agent, an inhibitor and/or promoter of cellular adhesion and/or growth (e.g. somatostatin analogues like lanreotide or octreotide, glucocorticoids, glycerol, heparin, botulinum toxin, a cytotoxic agent, a cytostatic agent, etc.) into the appropriate artery locations and inflating it to release said drug, agent, inhibitor or promoter;
- introducing the invention balloon coated with an anti-ulcer drug (e.g. omeprazole, esomeprazole or the like) into the stomach and inflating it to release said anti-ulcer drug; or
- introducing the invention balloon coated with an anti-inflammation drug into the colon, bladder or bowel and inflating it to release said anti-inflammation drug.

Of course, the various general improvements indicated for botulinum toxin formulations to be used on balloons according to this invention (such as, among many other things, the use of penetration enhancers) apply *mutatis mutandis* to the corresponding other drug formulations.

- 20 The term "about" refers to an interval around the considered value. As used in this patent application, "about X" means an interval from X minus 10% of X to X plus 10% of X, and preferably an interval from X minus 5% of X to X plus 5% of X.

- 25 Unless they are defined differently, all the technical and scientific terms used here have the same meaning as that usually understood by an ordinary specialist in the field to which this invention belongs. Similarly, all publications, patent applications, all patents and all other references mentioned here are incorporated by way of reference.

The following example is presented to illustrate the above and must in no case be considered as a limit to the scope of the invention.

**EXAMPLE**

A woman in her sixties suffers from urinary incontinence due to unstable bladder. She is infused in the bladder, through a catheter introduced into the urethra and the bladder and with the assistance of an endoscope apparatus, with 50 ml of a physiological saline solution containing 2,000 units of botulinum toxin type A (Dysport®; supplier: Ipsen Ltd, Wrexham, UK).

**Pharmacological procedures*****Bladder function***

The bladder capacity (in ml) is measured in patients before and after treatment at regular time intervals for up to 12 months. Additionally voiding function is recorded over this period.

Claims

1. Use of a solid, semi-solid or liquid formulation of botulinum toxin for the preparation of a medicament intended to treat a disorder characterised by bladder spasms wherein said medicament is for administration without using an injection into the bladder wall.
- 5 2. Use according to claim 1, wherein the disorder characterised by bladder spasms is selected from the group consisting of urinary incontinence due to unstable bladder or unstable detrusor sphincter, voiding complications due to detrusor overactivity or unstable detrusor sphincter, urinary retention secondary to spastic sphincter or hypertrophied bladder neck and neurogenic bladder dysfunction secondary to  
10 Parkinson's disease, spinal cord injury, stroke or multiple sclerosis or characterised by a spasm reflex.
3. Use according to claim 1 or 2, wherein the medicament prepared is in the form of a liquid or semi-solid formulation intended to be administered through infusion into the  
15 bladder or a solid, a gel formulation intended to be deposited at the appropriate location of the patient's bladder, a spray formulation intended to be administered through spraying at the appropriate location of the patient's bladder or a solid, semi-solid or liquid botulinum toxin formulation spread on the outer wall of a balloon intended to be inflated inside the bladder.
4. Use according to claim 3, wherein the medicament prepared is a liquid or  
20 semi-solid formulation for administration through infusion into the bladder.
5. Use according to claim 4, wherein the liquid or semi-solid formulation has a volume of 20 to 80 ml.
6. Use according to claim 3, wherein the liquid or semi-solid formulation contains from 100 to 2500 units of botulinum toxin type A.
- 25 7. Use according to claim 3, wherein the liquid or semi-solid formulation contains from 4,000 to 50,000 units of botulinum toxin type B.
8. Use according to claim 3, wherein the medicament prepared is in the form of a gel formulation.
9. Use according to claim 3, wherein the medicament prepared is in the form of a spray  
30 formulation.

C/SCRAS SPECS/46978,GB01.Spec

4 December 2003



10. Use according to claim 3, wherein the medicament prepared is in the form of a solid, semi-solid or liquid botulinum toxin formulation spread on the outer wall of a balloon intended to be inflated inside the bladder.

- 5 11. A method of treatment of a disorder characterised by bladder spasms, wherein the disorder characterised by bladder spasms is selected from the group consisting of urinary incontinence due to unstable bladder or unstable detrusor sphincter, voiding complications due to detrusor overactivity or unstable detrusor sphincter, urinary retention secondary to spastic sphincter or hypertrophied bladder neck and neurogenic bladder dysfunction secondary to Parkinson's disease, spinal cord injury, stroke or
- 10 multiple sclerosis or characterised by a spasm reflex: wherein a patient requiring said treatment is administered with a therapeutically effective dose of a solid, semi-solid or liquid formulation of botulinum toxin, wherein said formulation is administered without using an injection into the bladder wall.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**